

**Data Validation (DV) Report**  
**for TESS ID 283722336**  
**Sectors 17 - 17**

This Data Validation Report was produced in the  
TESS Science Processing Operations Center (SPOC) Pipeline  
at NASA Ames Research Center

15-Nov-2019 03:17:52 Z

# Contents

<b>1</b>	<b>Summary</b>	<b>1</b>
<b>2</b>	<b>Survey Image</b>	<b>2</b>
<b>3</b>	<b>Flux Time Series</b>	<b>3</b>
<b>4</b>	<b>Dashboards</b>	<b>5</b>
<b>5</b>	<b>Pixel Level Diagnostics</b>	<b>7</b>
5.1	Planet Candidate 1 . . . . .	7
5.2	Planet Candidate 2 . . . . .	11
5.3	Difference Image TIC Key . . . . .	15
<b>6</b>	<b>Phased Light Curves</b>	<b>16</b>
<b>7</b>	<b>Planet Candidate 1</b>	<b>20</b>
7.1	Model Fitter: All Transits . . . . .	20
7.2	Model Fitter: Reduced Parameter Fit Results . . . . .	23
7.3	Model Fitter: Trapezoidal Fit Results . . . . .	25
7.4	Validation Tests . . . . .	27
7.4.1	Weak Secondary Test . . . . .	27
7.4.2	Eclipsing Binary Discrimination Test . . . . .	27
7.4.3	Bootstrap Test . . . . .	28
7.4.4	Ghost Diagnostic Test . . . . .	28
7.4.5	Validation Test Figures . . . . .	29
<b>8</b>	<b>Planet Candidate 2</b>	<b>32</b>
8.1	Model Fitter: All Transits . . . . .	32
8.2	Model Fitter: Reduced Parameter Fit Results . . . . .	35
8.3	Model Fitter: Trapezoidal Fit Results . . . . .	37
8.4	Validation Tests . . . . .	39
8.4.1	Weak Secondary Test . . . . .	39
8.4.2	Eclipsing Binary Discrimination Test . . . . .	39
8.4.3	Bootstrap Test . . . . .	40
8.4.4	Ghost Diagnostic Test . . . . .	40
8.4.5	Validation Test Figures . . . . .	41
<b>Appendices</b>		<b>44</b>
<b>A</b>	<b>Planet Candidate 1</b>	<b>44</b>
A.1	Model Fitter: All Transits . . . . .	44
A.2	Model Fitter: Odd & Even Transits . . . . .	46
A.3	Eclipsing Binary Discrimination Test . . . . .	51
<b>B</b>	<b>Planet Candidate 2</b>	<b>52</b>
B.1	Model Fitter: All Transits . . . . .	52

B.2 Model Fitter: Odd & Even Transits . . . . .	54
B.3 Eclipsing Binary Discrimination Test . . . . .	59
<b>C Alerts</b>	<b>60</b>


# 1 Summary

Target Properties	Value	Uncertainty	Units	Provenance
Catalog ID	283722336			
TOI ID	-			
TESS Name	-			
RA	348.32072824	0	degrees	TIC8
Dec	57.16835459	0	degrees	TIC8
Magnitude	4.6278	0.0065		TIC8
Radius	0.751	0.058	Solar radii	TIC8
Effective Temperature	4884	143	Kelvin	TIC8
log(g)	4.590	0.099051	cm/sec <sup>2</sup>	TIC8
[M/H]	0.096	0.020813	Solar metallicity	TIC8
Stellar Density	1.892	0.455	Solar density	TIC8-Derived
Limb Darkening Coefficient 1	0.69802			
Limb Darkening Coefficient 2	-0.59897			
Limb Darkening Coefficient 3	1.1059			
Limb Darkening Coefficient 4	-0.46573			
Number of Planet Candidates	2			
TOI Model	toi-plus-2019-11-12_edited-2.csv			
TESS Names Model	-			
External TCE Model	-			
Software Revision	spoc-4.0.11-20191024			
Date Report Generated	15-Nov-2019 03:17:52 Z			

Sector	Target Table	Camera/ CCD	Crowding Metric	Flux Fraction
17	176	3:1	0.9979	0.9534


Planet Candidate	TOI ID	TESS Name	TOI Correlation	Period (days)	Period Ratio	Epoch (BTJD)	Semi-major Axis (AU)	Radius (Re)	Seff	Teq (K)	False Alarm	Suspected EB
1	-	-	-	3.093	1.00	1765.955	0.04	1.3	192.9	951	3.32e-21	false
2	-	-	-	6.765	2.19	1766.169	0.07	1.5	67.9	732	4.83e-22	false

## 2 Survey Image




Digitized Sky Survey (DSS) red image. The 5' x 5' image is centered on the J2000 coordinates of target (283722336).

### 3 Flux Time Series



Summary plot of sector-stitched flux time series and transits for target 283722336, marked with DV fitted epoch/period (or TPS epoch/period if fit was not successful). Transits of identified planets are labeled with epoch BTJD and orbital period. For the data of sector 17, target table 176, start BJD is 2458764. Open `./summary-plots/0000000283722336-00-flux-dv-fit-17-176.fig`



Summary plot of raw flux time series. For the data of sector 17, target table 176, start BJD is 2458764.

Open `./summary-plots/000000283722336-00-raw-flux-17-176.fig`

## 4 Dashboards

## Planet Candidate 1

<b>Model Fitter</b>	<b>Stellar Radius</b> 0.8 ± 0.1 Solar units		<b>Core Aperture Correlation Statistic</b> Value = <i>N/A</i> Significance = <i>N/A</i>		<b>Ghost Diagnostic Test</b>
	Period = 3.1 ± 0.0 days Depth = 271 ± 16 ppm Planet Radius = 1.3 ± 0.6 Earth radii Semi-major Axis = 0.0 ± 0.0 AU Effective Stellar Flux = 192.9 ± 38.3 Equilibrium Temperature = 951 ± 47 Kelvin Chi-squared/DoF = 0.9 SNR = 18.3		<b>Halo Aperture Correlation Statistic</b> Value = <i>N/A</i> Significance = <i>N/A</i>  <b>Core/Halo Ratio</b> Ratio = <i>N/A</i>		
<b>Eclipsing Binary Discrimination Test</b>	<b>Odd-Even Depth Comparison Statistic</b> Value = 1.78e-03 Significance = 96.63%		<b>Offsets Relative to Out of Transit Centroid</b> Source RA Offset = 2.93e+01 ± 2.51e+00 arcsec (11.67 $\sigma$ ) Source Dec Offset = 3.45e+00 ± 2.50e+00 arcsec (1.38 $\sigma$ ) Source Offset Distance = 2.95e+01 ± 2.51e+00 arcsec (11.75 $\sigma$ )  <b>Offsets Relative to TIC Position</b> Source RA Offset = 2.93e+01 ± 2.50e+00 arcsec (11.69 $\sigma$ ) Source Dec Offset = -2.62e+00 ± 2.50e+00 arcsec (-1.05 $\sigma$ ) Source Offset Distance = 2.94e+01 ± 2.50e+00 arcsec (11.74 $\sigma$ )		<b>Difference Image Centroid Offsets</b>
	<b>Shorter Period Comparison Statistic</b> Value = <i>N/A</i> Significance = <i>N/A</i>	<b>Longer Period Comparison Statistic</b> Value = 2.30e+03 Significance = 100.00%	False Alarm = 3.32e-21 Transit Count = 8 Max Multiple Event Statistic = 15.8		

Summary of model fitter results and validation test results for target 283722336, planet candidate 1. In general, green denotes that the candidate is likely a planet, while red denotes that the candidate is unlikely to be a planet. Cyan denotes that no data is available. The color of the Model Fitter block is: green, when the SNR of the fit is greater than or equal to 10; yellow, if the SNR is greater than or equal to 7.1 but less than 10; red, if the SNR is less than 7.1 or if the fitter failed. The color of the Ghost Diagnostic Test and Eclipsing Binary Discrimination Test blocks are: green, when the significance is within 2-sigma; yellow, when the significance is between 2- and 3-sigma; red when the significance is greater than 3-sigma. The color of the Difference Image Centroid Offsets block is: green, when the max offset distance sigma is less than or equal to 2; yellow, when the max sigma is between 2 and 3; red when the max sigma is greater than 3. The color of the Bootstrap Test block is green whenever the false alarm probability is less than  $10^{-12}$ , low enough to limit the total number of false alarms from a four year mission to less than one. If the false alarm probability is greater than  $10^{-12}$ , the color of the Bootstrap Test block is: green, when the false alarm probability is less than or equal to the CCDF of a Gaussian distribution at the observed maximum multiple event statistic; yellow when the false alarm probability is between 1 and 2 times that of a Gaussian distribution at the max multiple event statistic; and red when the false alarm probability is more than 2 times that of a Gaussian distribution at the max multiple event statistic.



## Planet Candidate 2

<b>Model Fitter</b>	<b>Stellar Radius</b> 0.8 ± 0.1 Solar units		<b>Core Aperture Correlation Statistic</b> Value = <i>N/A</i> Significance = <i>N/A</i>	<b>Ghost Diagnostic Test</b>
	Period = 6.8 ± 0.0 days Depth = 284 ± 16 ppm Planet Radius = 1.5 ± 0.2 Earth radii Semi-major Axis = 0.1 ± 0.0 AU Effective Stellar Flux = 67.9 ± 13.5 Equilibrium Temperature = 732 ± 36 Kelvin Chi-squared/DoF = 0.7 SNR = 16.9		<b>Halo Aperture Correlation Statistic</b> Value = <i>N/A</i> Significance = <i>N/A</i>  <b>Core/Halo Ratio</b> Ratio = <i>N/A</i>	
<b>Eclipsing Binary Discrimination Test</b>	<b>Odd-Even Depth Comparison Statistic</b> Value = 6.55e-01 Significance = 41.82%		<b>Offsets Relative to Out of Transit Centroid</b> Source RA Offset = -1.43e+02 ± 7.33e+00 arcsec (-19.52 $\sigma$ ) Source Dec Offset = -8.46e+01 ± 1.34e+01 arcsec (-6.30 $\sigma$ ) Source Offset Distance = 1.66e+02 ± 9.30e+00 arcsec (17.87 $\sigma$ )  <b>Offsets Relative to TIC Position</b> Source RA Offset = -1.43e+02 ± 7.24e+00 arcsec (-19.78 $\sigma$ ) Source Dec Offset = -9.07e+01 ± 1.34e+01 arcsec (-6.76 $\sigma$ ) Source Offset Distance = 1.69e+02 ± 9.43e+00 arcsec (17.96 $\sigma$ )	<b>Difference Image Centroid Offsets</b>
	<b>Shorter Period Comparison Statistic</b> Value = 2.30e+03 Significance = 100.00%	<b>Longer Period Comparison Statistic</b> Value = <i>N/A</i> Significance = <i>N/A</i>	False Alarm = 4.83e-22 Transit Count = 4 Max Multiple Event Statistic = 16.3	<b>Bootstrap Test</b>

Summary of model fitter results and validation test results for target 283722336, planet candidate 2. In general, green denotes that the candidate is likely a planet, while red denotes that the candidate is unlikely to be a planet. Cyan denotes that no data is available. The color of the Model Fitter block is: green, when the SNR of the fit is greater than or equal to 10; yellow, if the SNR is greater than or equal to 7.1 but less than 10; red, if the SNR is less than 7.1 or if the fitter failed. The color of the Ghost Diagnostic Test and Eclipsing Binary Discrimination Test blocks are: green, when the significance is within 2-sigma; yellow, when the significance is between 2- and 3-sigma; red when the significance is greater than 3-sigma. The color of the Difference Image Centroid Offsets block is: green, when the max offset distance sigma is less than or equal to 2; yellow, when the max sigma is between 2 and 3; red when the max sigma is greater than 3. The color of the Bootstrap Test block is green whenever the false alarm probability is less than  $10^{-12}$ , low enough to limit the total number of false alarms from a four year mission to less than one. If the false alarm probability is greater than  $10^{-12}$ , the color of the Bootstrap Test block is: green, when the false alarm probability is less than or equal to the CCDF of a Gaussian distribution at the observed maximum multiple event statistic; yellow when the false alarm probability is between 1 and 2 times that of a Gaussian distribution at the max multiple event statistic; and red when the false alarm probability is more than 2 times that of a Gaussian distribution at the max multiple event statistic.

## 5 Pixel Level Diagnostics

To reduce clutter, the catalog IDs in the difference images have been replaced by indices representing distance from the target star. The mapping between the indices and the catalog IDs is found in a table at the end of this section.

### 5.1 Planet Candidate 1

#### Multi-Sector Average PRF Fit of the Difference Images



Mean offset from the PRF fit to the out of transit image

	RA	Dec	Units
Offset	$29.3340 \pm 2.51e + 00$	$3.4467 \pm 2.50e + 00$	arcseconds
Offset/ $\sigma$	11.67	1.38	
Offset Distance	$29.5358 \pm 2.51e + 00$		arcseconds
Offset Distance/ $\sigma$	11.75		
$3\sigma$ Radius	7.5422		arcseconds

Mean offset from the TIC RA and Dec

	RA	Dec	Units
Offset	$29.2625 \pm 2.50e + 00$	$-2.6193 \pm 2.50e + 00$	arcseconds
Offset/ $\sigma$	11.69	-1.05	
Offset Distance	$29.3795 \pm 2.50e + 00$		arcseconds
Offset Distance/ $\sigma$	11.74		
$3\sigma$ Radius	7.5085		arcseconds



#### Planet Candidate 1



Difference image centroid offsets for target 283722336, planet candidate 1. Left: difference image PRF centroid offsets in RA and Dec with respect to the per sector out-of-transit centroids for the given target. Right: difference image PRF centroid offsets in RA and Dec with respect to the TC coordinates of the given target. Symbol key: green cross: per sector centroid offsets with 1-sigma error bars in RA and Dec; magenta cross: robust weighted mean offset over all sectors with 1-sigma error bars in RA and Dec; blue circle: 3-sigma radius of confusion for weighted mean offset; red asterisk: location of target star; blue asterisk: location of other TIC objects in the neighborhood. TIC ID and magnitude are noted in the text associated with each marked object. A constant error term of 2.5000 arcseconds has been added in quadrature to the computed uncertainty in the RA and Dec components of the robust mean offset.

Open `./planet-01/difference-image/0000000283722336-01-difference-image-centroid-offsets.fig`

## Planet Candidate 1




Difference image centroid offsets for target 283722336, planet candidate 1, displayed on survey image for given target. Left: difference image PRF centroid offsets in RA and Dec with respect to the per sector out-of-transit centroids for the given target. Right: difference image PRF centroid offsets in RA and Dec with respect to the TIC coordinates of the given target. Symbol key: green cross: per sector centroid offsets with 1-sigma error bars in RA and Dec; magenta cross: robust weighted mean offset over all sectors with 1-sigma error bars in RA and Dec; blue circle: 3-sigma radius of confusion for weighted mean offset; red asterisk: location of target star; blue asterisk: location of other TIC objects in the neighborhood. TIC ID and magnitude are noted in the text associated with each marked object. A constant error term of 2.5000 arcseconds has been added in quadrature to the computed uncertainty in the RA and Dec components of the robust mean offset.

Open `./planet-01/difference-image/0000000283722336-01-difference-image-centroid-offsets-survey.fig`

## Difference Image Summary Metrics

Number of Difference Images	Number of Metrics	Number of Good Metrics	Fraction of Good Metrics	Quality Threshold
1	1	0	0.0000	0.70

**Difference Image**  
Planet Candidate 1 / Sector 17 / Target Pixel Table 176



Difference image for target 283722336, planet candidate 1, sector 17, target pixel table 176. Upper left: difference between mean flux out-of-transit and in-transit; upper right: mean out-of-transit flux; lower left: mean in-transit flux; lower right: difference between mean flux out-of-transit and in-transit after normalizing by the uncertainty in the difference for each pixel. The optimal aperture is outlined with a white dash-dotted line in each panel and the target mask is outlined with a solid white line. Symbol key: x: target position from TIC RA and Dec converted to CCD coordinates via motion polynomials; \*: position of nearby TIC objects converted to CCD coordinates via motion polynomials; +: PRF-fit location of target from out-of-transit image; triangle: PRF-fit location of transit source from the difference image. Number of transits = 8; number of valid in-transit cadences = 168; number of in-transit cadence gaps = 0; number of valid out-of-transit cadences = 480; number of out-of-transit cadence gaps = 0. Difference image quality metric = 0.44 (not good).

Open `./planet-01/difference-image/0000000283722336-01-difference-image-17-176.fig`

## PRF Fit of the Difference Image

## Offset from the PRF fit to the out of transit image

	Row	Column	Units	RA	Dec	Units
Out of Transit Image Centroid	$105.64 \pm 4.40e - 06$	$1869.54 \pm 4.67e - 06$	pixels	$348.34173767 \pm 1.55e - 06$	$57.16829178 \pm 1.46e - 06$	degrees
Difference Image Centroid	$104.72 \pm 9.87e - 03$	$1870.67 \pm 6.57e - 03$	pixels	$348.35676670 \pm 6.09e - 05$	$57.16924921 \pm 3.08e - 05$	degrees
Offset	$-0.9167 \pm 9.87e - 03$	$1.1283 \pm 6.57e - 03$	pixels	$29.3340 \pm 2.67e - 01$	$3.4467 \pm 1.11e - 01$	arcseconds
Offset/ $\sigma$	-92.84	171.71		109.77	31.08	
Offset Distance	$1.4537 \pm 9.85e - 03$		pixels	$29.5358 \pm 2.65e - 01$		arcseconds
Offset Distance/ $\sigma$	147.58			111.44		

## Offset from the TIC RA and Dec converted to pixels via motion polynomials



	Row	Column	Units	RA	Dec	Units
TIC Reference Centroid	$105.85 \pm 2.75e - 04$	$1869.76 \pm 2.50e - 04$	pixels	$348.34177363 \pm 0.00e + 00$	$57.16997679 \pm 0.00e + 00$	degrees
Difference Image Centroid	$104.72 \pm 9.87e - 03$	$1870.67 \pm 6.57e - 03$	pixels	$348.35676670 \pm 6.09e - 05$	$57.16924921 \pm 3.08e - 05$	degrees
Offset	$-1.1342 \pm 9.88e - 03$	$0.9166 \pm 6.58e - 03$	pixels	$29.2625 \pm 1.19e - 01$	$-2.6193 \pm 1.11e - 01$	arcseconds
Offset/ $\sigma$	-114.83	139.40		246.02	-23.64	
Offset Distance	$1.4583 \pm 1.04e - 02$		pixels	$29.3795 \pm 1.21e - 01$		arcseconds
Offset Distance/ $\sigma$	140.11			242.90		

## 5.2 Planet Candidate 2

## Multi-Sector Average PRF Fit of the Difference Images

Mean offset from the PRF fit to the out of transit image				Mean offset from the TIC RA and Dec			
	RA	Dec	Units		RA	Dec	Units
Offset	$-143.0570 \pm 7.33e + 00$	$-84.6135 \pm 1.34e + 01$	arcseconds	Offset	$-143.1110 \pm 7.24e + 00$	$-90.6912 \pm 1.34e + 01$	arcseconds
Offset/ $\sigma$	-19.52	-6.30		Offset/ $\sigma$	-19.78	-6.76	
Offset Distance	$166.2070 \pm 9.30e + 00$		arcseconds	Offset Distance	$169.4274 \pm 9.43e + 00$		arcseconds
Offset Distance/ $\sigma$	17.87			Offset Distance/ $\sigma$	17.96		
$3\sigma$ Radius	27.8979		arcseconds	$3\sigma$ Radius	28.2970		arcseconds



## Planet Candidate 2



Difference image centroid offsets for target 283722336, planet candidate 2. Left: difference image PRF centroid offsets in RA and Dec with respect to the per sector out-of-transit centroids for the given target. Right: difference image PRF centroid offsets in RA and Dec with respect to the TC coordinates of the given target. Symbol key: green cross: per sector centroid offsets with 1-sigma error bars in RA and Dec; magenta cross: robust weighted mean offset over all sectors with 1-sigma error bars in RA and Dec; blue circle: 3-sigma radius of confusion for weighted mean offset; red asterisk: location of target star; blue asterisk: location of other TIC objects in the neighborhood. TIC ID and magnitude are noted in the text associated with each marked object. A constant error term of 2.5000 arcseconds has been added in quadrature to the computed uncertainty in the RA and Dec components of the robust mean offset.

Open `./planet-02/difference-image/0000000283722336-02-difference-image-centroid-offsets.fig`

## Planet Candidate 2




Difference image centroid offsets for target 283722336, planet candidate 2, displayed on survey image for given target. Left: difference image PRF centroid offsets in RA and Dec with respect to the per sector out-of-transit centroids for the given target. Right: difference image PRF centroid offsets in RA and Dec with respect to the TIC coordinates of the given target. Symbol key: green cross: per sector centroid offsets with 1-sigma error bars in RA and Dec; magenta cross: robust weighted mean offset over all sectors with 1-sigma error bars in RA and Dec; blue circle: 3-sigma radius of confusion for weighted mean offset; red asterisk: location of target star; blue asterisk: location of other TIC objects in the neighborhood. TIC ID and magnitude are noted in the text associated with each marked object. A constant error term of 2.5000 arcseconds has been added in quadrature to the computed uncertainty in the RA and Dec components of the robust mean offset.

Open `./planet-02/difference-image/0000000283722336-02-difference-image-centroid-offsets-survey.fig`

## Difference Image Summary Metrics

Number of Difference Images	Number of Metrics	Number of Good Metrics	Fraction of Good Metrics	Quality Threshold
1	1	0	0.0000	0.70

**Difference Image**  
Planet Candidate 2 / Sector 17 / Target Pixel Table 176



Difference image for target 283722336, planet candidate 2, sector 17, target pixel table 176. Upper left: difference between mean flux out-of-transit and in-transit; upper right: mean out-of-transit flux; lower left: mean in-transit flux; lower right: difference between mean flux out-of-transit and in-transit after normalizing by the uncertainty in the difference for each pixel. The optimal aperture is outlined with a white dash-dotted line in each panel and the target mask is outlined with a solid white line. Symbol key: x: target position from TIC RA and Dec converted to CCD coordinates via motion polynomials; \*: position of nearby TIC objects converted to CCD coordinates via motion polynomials; +: PRF-fit location of target from out-of-transit image; triangle: PRF-fit location of transit source. Number of transits = 4; number of valid in-transit cadences = 160; number of in-transit cadence gaps = 0; number of valid out-of-transit cadences = 421; number of out-of-transit cadence gaps = 0. Difference image quality metric = -0.00 (not good).

Open `./planet-02/difference-image/0000000283722336-02-difference-image-17-176.fig`



## PRF Fit of the Difference Image

## Offset from the PRF fit to the out of transit image

	Row	Column	Units	RA	Dec	Units
Out of Transit Image Centroid	$105.64 \pm 4.71e - 06$	$1869.54 \pm 4.98e - 06$	pixels	$348.34174265 \pm 1.54e - 06$	$57.16828854 \pm 1.46e - 06$	degrees
Difference Image Centroid	$107.66 \pm 5.45e - 01$	$1861.70 \pm 6.94e - 01$	pixels	$348.26844862 \pm 3.48e - 03$	$57.14478478 \pm 3.66e - 03$	degrees
Offset	$2.0179 \pm 5.45e - 01$	$-7.8411 \pm 6.94e - 01$	pixels	$-143.0570 \pm 6.89e + 00$	$-84.6135 \pm 1.32e + 01$	arcseconds
Offset/ $\sigma$	3.70	-11.29		-20.76	-6.42	
Offset Distance	$8.0966 \pm 6.88e - 01$		pixels	$166.2070 \pm 9.94e + 00$		arcseconds
Offset Distance/ $\sigma$	11.76			16.73		

## Offset from the TIC RA and Dec converted to pixels via motion polynomials


	Row	Column	Units	RA	Dec	Units
TIC Reference Centroid	$105.86 \pm 2.74e - 04$	$1869.76 \pm 2.49e - 04$	pixels	$348.34177363 \pm 0.00e + 00$	$57.16997679 \pm 0.00e + 00$	degrees
Difference Image Centroid	$107.66 \pm 5.45e - 01$	$1861.70 \pm 6.94e - 01$	pixels	$348.26844862 \pm 3.48e - 03$	$57.14478478 \pm 3.66e - 03$	degrees
Offset	$1.7997 \pm 5.45e - 01$	$-8.0529 \pm 6.94e - 01$	pixels	$-143.1110 \pm 6.79e + 00$	$-90.6912 \pm 1.32e + 01$	arcseconds
Offset/ $\sigma$	3.30	-11.60		-21.07	-6.88	
Offset Distance	$8.2515 \pm 6.90e - 01$		pixels	$169.4274 \pm 1.01e + 01$		arcseconds
Offset Distance/ $\sigma$	11.96			16.79		

### 5.3 Difference Image TIC Key

Index	Catalog ID	Mag	RA (degrees)	Dec (degrees)	Distance (arcsec)
1	283722336	4.628	348.34177363	57.16997679	0.00
2	283722382	13.009	348.36318837	57.17811194	51.04
3	283722412	13.042	348.33758368	57.18451062	52.96
4	2047267705	13.464	348.33760991	57.18452821	53.01
5	283722251	12.540	348.36895582	57.15138951	85.39
6	283722394	10.994	348.29656324	57.17998879	95.32


RA, Dec and Distances are corrected for proper motion. This table may not contain all of the objects shown.

## 6 Phased Light Curves



Phased unwhitened flux time series is plotted in black dots. When all transits fit completed with full or secondary convergence, the phase is determined with the fitted epoch and period; otherwise, the phase is determined with the TPS epoch and period. The values of the phased unwhitened flux time series averaged in one cadence wide bins are plotted in bigger blue dots. When all transits fit completes with full or secondary convergence, the averaged values of the phased unwhitened fitted model light curve are plotted in red dots. Transit event markers in different colors indicate the locations of the transits of all planet candidates. The transits of the same planet candidate are labeled with the markers of the same color, for example, blue markers for transits of plane candidate #1, red markers for transits of planet candidate #2, etc.


Open `./summary-plots/0000000283722336-01-phased-unwhitened-flux-time-series.fig`



Phased whitened flux time series is plotted in black dots. When all transits fit completed with full or secondary convergence, the phase is determined with the fitted epoch and period; otherwise, the phase is determined with the TPS epoch and period. The values of the phased whitened flux time series averaged in one cadence wide bins are plotted in bigger blue dots. When all transits fit completes with full or secondary convergence, the averaged values of the phased whitened fitted model light curve are plotted in red dots. Transit event markers in different colors indicate the locations of the transits of all planet candidates. The transits of the same planet candidate are labeled with the markers of the same color, for example, blue markers for transits of plane candidate #1, red markers for transits of planet candidate #2, etc.


Open `./summary-plots/0000000283722336-01-phased-whitened-flux-time-series.fig`

## Planet: 1 Phased Unwhitened Flux Time Series by Sector



Phased unwhitened flux time series by sector for target 283722336, planet candidate 1. Period = 3.0931 days; transit epoch = 1765.9545 BTJD.  
Open `./summary-plots/000000283722336-01-phased-unwhitened-flux-time-series-by-sector.fig`

## Planet: 2 Phased Unwhitened Flux Time Series by Sector



Phased unwhitened flux time series by sector for target 283722336, planet candidate 2. Period = 6.7654 days; transit epoch = 1766.1693 BTJD.  
Open `./summary-plots/000000283722336-02-phased-unwhitened-flux-time-series-by-sector.fig`

## 7 Planet Candidate 1

### 7.1 Model Fitter: All Transits


Model Characteristic	Name
Transit Model	mandel-agol_geometric_transit_model
Limb Darkening Model	claret_tess_nonlinear_limb_darkening_model

TCE Parameter	Value	Units
Trial Transit Pulse Duration	1.0	hours
Transit Epoch	1765.9518053	TJD
Orbital Period	3.0931370	days
Maximum SES	7.3	
Maximum MES	15.8	
Robust Statistic	15.0	
Chi Square Goodness of Fit Statistic (DoF)	399.4 (238)	
Chi Square2 Statistic (DoF)	3.6 (29.1)	
Threshold for Desired PFA		

DoF: Degrees of Freedom

Parameter	Value	Uncertainty	Units
SNR	18.3		
Orbital Period	3.0930660	2.3922e-04	days
Transit Epoch	1765.9545257	1.0484e-03	BTJD
Impact Parameter	0.5983	1.9588e+00	
Planet Radius to Star Radius Ratio	0.0157252	6.6071e-03	
Semi-major Axis to Star Radius Ratio	22.5905	4.0338e+01	
Planet Radius	1.2886	5.5034e-01	Earth radii
Semi-major Axis	0.0386	3.5349e-03	AU
Effective Stellar Flux	192.9197	3.8300e+01	Goldilocks
Equilibrium Temperature	951	4.7176e+01	Kelvin
Stellar Density	16.1896	8.6724e+01	Solar density
Transit Depth	271	1.5627e+01	ppm
Transit Duration	0.8591	5.7487e-02	hours
Transit Ingress Duration	0.0205	8.2919e-02	hours
Eccentricity	0.0000	0.0000e+00	
Peri Longitude	0.0000	0.0000e+00	degrees
Model Chi Square Statistic (DoF)	1016.3 (1102.2)		
Model Chi Square Goodness of Fit Statistic (DoF)	151.1 (231)		
Model Chi Square2 Statistic (DoF)	4.5 (7)		


DoF: Degrees of Freedom



Flux time series for CatId 283722336, Planet candidate 1 in the unwhitened domain. For the data of Sector-17/TargetTableId-176, start BJD is 2458764. Transit event markers indicate the location of transits of the given planet candidate. All transits fit completed with full convergence.


Open `./planet-01/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-01-all-unwhitened-17-176.fig`





Folded flux time series for CatId 283722336, Planet candidate 1 in the whitenened domain is plotted in black dots. Values are averaged into 1 cadence wide bins. The blue dots represent the averaged values of the folded flux time series; the red dots represent the averaged values of the folded model light curve of the all transits fit; the green dots are the averaged folded fit residuals, vertically offset for clarity. All transits fit completed with full convergence.

Open `./planet-01/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-01-all-whitenened.fig`




Folded flux time series for CatId 283722336, Planet candidate 1 in the whitenened domain, zoomed on the transit. The flux data whose robust weights are larger/smaller than 0.1 are plotted in dark green/cyan dots, respectively. Values are averaged into 1 cadence wide bins. The blue dots represent the averaged values of the folded flux time series; the red dots represent the averaged values of the fitted model light curve of the all transits fit; the green dots are the averaged folded fit residuals, vertically offset for clarity. Magenta dots are the averaged values of the folded flux time series, with a phase shift of 0.5 relative to the blue dots, vertically offset for clarity. All transits fit completed with full convergence.

Open `./planet-01/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-01-all-whitenened-zoomed.fig`

## 7.2 Model Fitter: Reduced Parameter Fit Results


Impact Parameter	SNR	Model Chi Square	Planet Radius to Star Radius	Uncert	Semi-major Axis to Star Radius	Uncert	Transit Depth (ppm)	Uncert	Transit Duration (hours)	Uncert
0.10	19.1	2075.7	0.0149524	4.8880e-04	27.4382	9.7243e-01	272	1.7682e+01	0.8699	3.0774e-02
0.30	19.1	2088.8	0.0150963	4.9597e-04	26.2379	9.3076e-01	272	1.7752e+01	0.8735	3.0927e-02
0.50	19.2	2055.6	0.0154510	5.0028e-04	23.9339	8.4194e-01	272	1.7500e+01	0.8729	3.0636e-02
0.70	19.3	2129.0	0.0161510	5.2961e-04	19.8280	7.0776e-01	271	1.7688e+01	0.8785	3.1265e-02
0.90	19.2	2092.9	0.0179255	5.8805e-04	12.3167	4.4497e-01	270	1.7630e+01	0.9150	3.2918e-02

Highlighted row is the best reduced-parameter model fit.




Model chi squares of reduced parameter fits vs. impact parameter for CatId 283722336, Planet candidate 1. The fit result with the minimum chi square is marked with a dashed line in the plot.

Open `./planet-01/planet-search-and-model-fitting-results/reduced-parameter-fits/0000000283722336-01-reduced-fits-chi-square.fig`



Ratios of planet radius to star radius of reduced parameter fits vs. impact parameter for CatId 283722336, Planet candidate 1. The fit result with the minimum chi square is marked with a dashed line in the plot.

Open `./planet-01/planet-search-and-model-fitting-results/reduced-parameter-fits/0000000283722336-01-reduced-fits-rp-over-rstar.fig`



Ratios of semimajor axis to star radius of reduced parameter fits vs. impact parameter for CatId 283722336, Planet candidate 1. The fit result with the minimum chi square is marked with a dashed line in the plot.

Open `./planet-01/planet-search-and-model-fitting-results/reduced-parameter-fits/0000000283722336-01-reduced-fits-a-over-rstar.fig`

### 7.3 Model Fitter: Trapezoidal Fit Results


Model Characteristic	Name
Transit Model	trapezoidal_model
Limb Darkening Model	

TCE Parameter	Value	Units
Trial Transit Pulse Duration	1.0	hours
Transit Epoch	1765.9518053	TJD
Orbital Period	3.0931370	days
Maximum SES	7.3	
Maximum MES	15.8	
Robust Statistic	15.0	
Chi Square Goodness of Fit Statistic (DoF)	399.4 (238)	
Chi Square2 Statistic (DoF)	3.6 (29.1)	
Threshold for Desired PFA		

DoF: Degrees of Freedom


Parameter	Value	Uncertainty	Units
SNR	1.6		
Orbital Period	3.0931370		days
Transit Epoch	1765.7699789		BTJD
Transit Depth	7		ppm
Transit Duration	2.7172		hours
Transit Ingress Duration	0.0060		hours
Model Chi Square Statistic (DoF)	17911.8 (1852)		

DoF: Degrees of Freedom



Folded detrended flux time series for CatId 283722336, Planet candidate 1 and folded trapezoidal model light curve.

Open `./planet-01/planet-search-and-model-fitting-results/trapezoidal-model-fit/0000000283722336-01-all-trapezoidal.fig`



Zoomed folded detrended flux time series for CatId 283722336, Planet candidate 1 and folded trapezoidal model light curve.

Open `./planet-01/planet-search-and-model-fitting-results/trapezoidal-model-fit/0000000283722336-01-all-trapezoidal-zoomed.fig`

## 7.4 Validation Tests

The Centroid Test and Eclipsing Binary Discrimination Test are chi-squared hypothesis tests. For these tests, a significance of 100% favors a planet, while 0% indicates an unlikely planet.

### 7.4.1 Weak Secondary Test

Result	Value	Uncertainty	Units	Statistic in Sigmas	Significance (%)
Orbital Period	3.0931		days		
Transit Duration	1		hours		
Maximum MES	15.8				
Secondary Phase	1.3639		days		
Secondary MES	4.7				
Minimum Phase	2.2306		days		
Minimum MES	-2.8				
Median MES	-0.1				
MAD MES	0.69112				
Robust Statistic	3.8				
Secondary Depth	71.3	2.2073e+01	ppm		
Geometric Albedo	35.2	3.2020e+01		1.0680	14.28
Planet Effective Temperature	3580	8.0816e+02	Kelvin	3.2476	0.06

### 7.4.2 Eclipsing Binary Discrimination Test

Result	Value	Value in Sigmas	Significance (%)
Odd Even Transit Depth Comparison Statistic	1.7835e-03	0.0422	96.63
Longer Period Comparison Statistic	2.2962e+03	47.9186	100.00


### 7.4.3 Bootstrap Test

Result	Value
False Alarm Probability	3.3156e-21
Bootstrap Threshold for Desired PFA	11.3
MES Mean	-2.92
MES Standard Deviation	1.99
Transit Count	8

### 7.4.4 Ghost Diagnostic Test

Result	Value	Significance (%)
Maximum MES	15.8	
SNR	18.3	
Core Aperture Statistic		
Halo Aperture Statistic		
Ratio of Core/Halo Aperture Statistics		


## 7.4.5 Validation Test Figures



The primary event has been set to zero and both the max and min of the resulting MES vs. Phase are marked with a red star. The best matched pulse duration in hours is 1. The maximum secondary MES and corresponding phase are 4.655 and 1.3639 days respectively. The minimum secondary MES and corresponding phase are -2.7972 and 2.2306 days respectively.

Open `./planet-01/report-summary/000000283722336-01-weak-secondary-diagnostic.fig`





Bootstrap results for target 283722336, planet 1. Cumulative sum of the probabilities (derived from the histogram of counts) from upper tail to the search transit threshold; false alarm probability is indicated by the star. The Gaussian equivalent threshold for this false alarm probability is 9.3795. The threshold on this distribution that achieves the same false alarm rate as a 7.1 sigma threshold on a Gaussian distribution is 11.3083.

Open `./planet-01/bootstrap-results/0000000283722336-01-bootstrap-false-alarm.fig`

No figures named 000000283722336-01-\*--unwhitened-cotrended-zoomed-model.fig are available.

## 8 Planet Candidate 2

### 8.1 Model Fitter: All Transits


Model Characteristic	Name
Transit Model	mandel-agol_geometric_transit_model
Limb Darkening Model	claret_tess_nonlinear_limb_darkening_model

TCE Parameter	Value	Units
Trial Transit Pulse Duration	1.5	hours
Transit Epoch	1766.1677774	TJD
Orbital Period	6.7652748	days
Maximum SES	8.4	
Maximum MES	16.3	
Robust Statistic	15.8	
Chi Square Goodness of Fit Statistic (DoF)	213.7 (178)	
Chi Square2 Statistic (DoF)	1.7 (25.7)	
Threshold for Desired PFA		

DoF: Degrees of Freedom


Parameter	Value	Uncertainty	Units
SNR	16.9		
Orbital Period	6.7654340	6.4710e-04	days
Transit Epoch	1766.1693222	1.2671e-03	BTJD
Impact Parameter	0.8691	1.6716e-01	
Planet Radius to Star Radius Ratio	0.0179089	2.2259e-03	
Semi-major Axis to Star Radius Ratio	16.8674	9.8382e+00	
Planet Radius	1.4675	2.1429e-01	Earth radii
Semi-major Axis	0.0650	5.9563e-03	AU
Effective Stellar Flux	67.9466	1.3489e+01	Goldilocks
Equilibrium Temperature	732	3.6343e+01	Kelvin
Stellar Density	1.4086	2.4648e+00	Solar density
Transit Depth	284	1.5872e+01	ppm
Transit Duration	1.6264	1.2855e-01	hours
Transit Ingress Duration	0.1113	1.4544e-01	hours
Eccentricity	0.0000	0.0000e+00	
Peri Longitude	0.0000	0.0000e+00	degrees
Model Chi Square Statistic (DoF)	679.7 (920.5)		
Model Chi Square Goodness of Fit Statistic (DoF)	92.8 (208)		
Model Chi Square2 Statistic (DoF)	0.5 (3)		

DoF: Degrees of Freedom




Flux time series for CatId 283722336, Planet candidate 2 in the unwhitened domain. For the data of Sector-17/TargetTableId-176, start BJD is 2458764. Transit event markers indicate the location of transits of the given planet candidate. All transits fit completed with full convergence.

Open `./planet-02/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-02-all-unwhitened-17-176.fig`



Folded flux time series for CatId 283722336, Planet candidate 2 in the whitened domain is plotted in black dots. Values are averaged into 1 cadence wide bins. The blue dots represent the averaged values of the folded flux time series; the red dots represent the averaged values of the folded model light curve of the all transits fit; the green dots are the averaged folded fit residuals, vertically offset for clarity. All transits fit completed with full convergence.

Open `./planet-02/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-02-all-whitened.fig`




Folded flux time series for CatId 283722336, Planet candidate 2 in the whitened domain, zoomed on the transit. The flux data whose robust weights are larger/smaller than 0.1 are plotted in dark green/cyan dots, respectively. Values are averaged into 1 cadence wide bins. The blue dots represent the averaged values of the folded flux time series; the red dots represent the averaged values of the fitted model light curve of the all transits fit; the green dots are the averaged folded fit residuals, vertically offset for clarity. Magenta dots are the averaged values of the folded flux time series, with a phase shift of 0.5 relative to the blue dots, vertically offset for clarity. All transits fit completed with full convergence.

Open `./planet-02/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-02-all-whitened-zoomed.fig`

## 8.2 Model Fitter: Reduced Parameter Fit Results


Impact Parameter	SNR	Model Chi Square	Planet Radius to Star Radius	Uncert	Semi-major Axis to Star Radius	Uncert	Transit Depth (ppm)	Uncert	Transit Duration (hours)	Uncert
0.10	17.2	1287.3	0.0150246	5.0648e-04	33.4601	9.2497e-01	275	1.8435e+01	1.5605	4.3068e-02
0.30	17.2	1284.0	0.0152136	5.1243e-04	32.7752	8.9686e-01	276	1.8510e+01	1.5298	4.1789e-02
0.50	17.3	1294.5	0.0155744	5.2330e-04	29.8372	8.1518e-01	276	1.8477e+01	1.5318	4.1775e-02
0.70	17.3	1294.8	0.0162675	5.4703e-04	24.5903	6.7531e-01	275	1.8427e+01	1.5495	4.2470e-02
0.90	17.3	1264.4	0.0180058	6.0364e-04	15.0572	4.4220e-01	273	1.8207e+01	1.6363	4.8211e-02

Highlighted row is the best reduced-parameter model fit.




Model chi squares of reduced parameter fits vs. impact parameter for CatId 283722336, Planet candidate 2. The fit result with the minimum chi square is marked with a dashed line in the plot.

Open `./planet-02/planet-search-and-model-fitting-results/reduced-parameter-fits/0000000283722336-02-reduced-fits-chi-square.fig`



Ratios of planet radius to star radius of reduced parameter fits vs. impact parameter for CatId 283722336, Planet candidate 2. The fit result with the minimum chi square is marked with a dashed line in the plot.

Open `./planet-02/planet-search-and-model-fitting-results/reduced-parameter-fits/0000000283722336-02-reduced-fits-rp-over-rstar.fig`



Ratios of semimajor axis to star radius of reduced parameter fits vs. impact parameter for CatId 283722336, Planet candidate 2. The fit result with the minimum chi square is marked with a dashed line in the plot.

Open `./planet-02/planet-search-and-model-fitting-results/reduced-parameter-fits/0000000283722336-02-reduced-fits-a-over-rstar.fig`

### 8.3 Model Fitter: Trapezoidal Fit Results

Model Characteristic	Name
Transit Model	trapezoidal_model
Limb Darkening Model	


TCE Parameter	Value	Units
Trial Transit Pulse Duration	1.5	hours
Transit Epoch	1766.1677774	TJD
Orbital Period	6.7652748	days
Maximum SES	8.4	
Maximum MES	16.3	
Robust Statistic	15.8	
Chi Square Goodness of Fit Statistic (DoF)	213.7 (178)	
Chi Square2 Statistic (DoF)	1.7 (25.7)	
Threshold for Desired PFA		

DoF: Degrees of Freedom

Parameter	Value	Uncertainty	Units
SNR	26.9		
Orbital Period	6.7652748		days
Transit Epoch	1766.1696298		BTJD
Transit Depth	271		ppm
Transit Duration	1.6536		hours
Transit Ingress Duration	0.1899		hours
Model Chi Square Statistic (DoF)	16934.0 (1350)		


DoF: Degrees of Freedom





Folded detrended flux time series for CatId 283722336, Planet candidate 2 and folded trapezoidal model light curve.

Open `./planet-02/planet-search-and-model-fitting-results/trapezoidal-model-fit/0000000283722336-02-all-trapezoidal.fig`



Zoomed folded detrended flux time series for CatId 283722336, Planet candidate 2 and folded trapezoidal model light curve.

Open `./planet-02/planet-search-and-model-fitting-results/trapezoidal-model-fit/0000000283722336-02-all-trapezoidal-zoomed.fig`

## 8.4 Validation Tests

The Centroid Test and Eclipsing Binary Discrimination Test are chi-squared hypothesis tests. For these tests, a significance of 100% favors a planet, while 0% indicates an unlikely planet.

### 8.4.1 Weak Secondary Test

Result	Value	Uncertainty	Units	Statistic in Sigmas	Significance (%)
Orbital Period	6.7653		days		
Transit Duration	1.5		hours		
Maximum MES	16.3				
Secondary Phase	-0.59722		days		
Secondary MES	2.7				
Minimum Phase	1.4097		days		
Minimum MES	-3.1				
Median MES	0.0				
MAD MES	0.61956				
Robust Statistic	0.7				
Secondary Depth	16.1	2.7059e+01	ppm		
Geometric Albedo	17.4	2.9678e+01		0.5539	28.98
Planet Effective Temperature	2314	9.8223e+02	Kelvin	1.6089	5.38

### 8.4.2 Eclipsing Binary Discrimination Test

Result	Value	Value in Sigmas	Significance (%)
Odd Even Transit Depth Comparison Statistic	6.5539e-01	0.8096	41.82
Shorter Period Comparison Statistic	2.2962e+03	47.9186	100.00


### 8.4.3 Bootstrap Test

Result	Value
False Alarm Probability	4.8272e-22
Bootstrap Threshold for Desired PFA	11.4
MES Mean	-2.59
MES Standard Deviation	1.97
Transit Count	4

### 8.4.4 Ghost Diagnostic Test


Result	Value	Significance (%)
Maximum MES	16.3	
SNR	16.9	
Core Aperture Statistic		
Halo Aperture Statistic		
Ratio of Core/Halo Aperture Statistics		

## 8.4.5 Validation Test Figures



The primary event has been set to zero and both the max and min of the resulting MES vs. Phase are marked with a red star. The best matched pulse duration in hours is 1.5. The maximum secondary MES and corresponding phase are 2.7365 and -0.59722 days respectively. The minimum secondary MES and corresponding phase are -3.069 and 1.4097 days respectively.

Open `./planet-02/report-summary/0000000283722336-02-weak-secondary-diagnostic.fig`




Bootstrap results for target 283722336, planet 2. Cumulative sum of the probabilities (derived from the histogram of counts) from upper tail to the search transit threshold; false alarm probability is indicated by the star. The Gaussian equivalent threshold for this false alarm probability is 9.5805. The threshold on this distribution that achieves the same false alarm rate as a 7.1 sigma threshold on a Gaussian distribution is 11.3841.

Open `./planet-02/bootstrap-results/0000000283722336-02-bootstrap-false-alarm.fig`

No figures named 000000283722336-02-\*-unwhitened-cotrended-zoomed-model.fig are available.


## Appendix A Planet Candidate 1

### A.1 Model Fitter: All Transits




Robust weights distribution for CatId 283722336, Planet candidate 1. Top plot: all data points. Middle plot: all data points, folded per the fitted period and epoch. Bottom plot: all data points, folded and zoomed.

Open `./planet-01/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-01-all-robust-weights.fig`



Fit residuals distribution for CatId 283722336, Planet candidate 1. Only the valid data points used to constrain the fit are shown here. A Gaussian fit to the histogram is shown in red.

Open `./planet-01/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-01-all-histo-used.fig`



Fit residuals distribution for CatId 283722336, Planet candidate 1. Top plot: all valid data. Bottom plot: valid data not used to constrain fit (due to distance from a transit). Gaussian fits to the histograms are shown in red.



Open `./planet-01/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-01-all-histo-all-and-unused.fig`



## A.2 Model Fitter: Odd &amp; Even Transits



Parameter	Odd Transits Value	Odd Transits Uncertainty	Even Transits Value	Even Transits Uncertainty	Units	$\frac{\text{Difference}}{\ \text{Uncertainty}\ }$
SNR	12.6		13.3			
Orbital Period	3.0929741	3.3957e-04	3.0931629	3.1957e-04	days	4.0487e-01
Transit Epoch	1765.9552519	1.3658e-03	1769.0469139	1.2586e-03	BTJD	7.5592e-01
Impact Parameter	0.8954	3.2721e-01	0.8074	1.1732e+00		7.2253e-02
Planet Radius to Star Radius Ratio	0.0178838	5.2524e-03	0.0169054	9.7468e-03		8.8373e-02
Semi-major Axis to Star Radius Ratio	13.2773	1.9118e+01	16.3803	4.3655e+01		6.5112e-02
Planet Radius	1.4654	4.4482e-01	1.3853	8.0571e-01	Earth radii	8.7117e-02
Semi-major Axis	0.0386	3.5348e-03	0.0386	3.5350e-03	AU	3.1410e-04
Effective Stellar Flux	192.9273	3.8301e+01	192.9116	3.8298e+01	Goldilocks	2.8986e-04
Equilibrium Temperature	951	4.7177e+01	951	4.7176e+01	Kelvin	2.8986e-04
Stellar Density	3.2871	1.4199e+01	6.1716	4.9344e+01	Solar density	5.6179e-02
Transit Depth	271	2.3567e+01	273	2.1917e+01	ppm	4.2232e-02
Transit Duration	0.8637	1.9457e-01	0.8932	2.0254e-01	hours	1.0485e-01
Transit Ingress Duration	0.0719	2.3299e-01	0.0414	2.4791e-01	hours	8.9562e-02
Eccentricity	0.0000	0.0000e+00	0.0000	0.0000e+00		
Peri Longitude	0.0000	0.0000e+00	0.0000	0.0000e+00	degrees	
Model Chi Square Statistic (DoF)	1017.9 (1097.8)		1017.9 (1097.8)			

DoF: Degrees of Freedom




Folded flux time series for CatId 283722336, Planet candidate 1 in the whitened domain is plotted in black dots. Values are averaged into 1 cadence wide bins. The blue dots represent the averaged values of the folded flux time series; the red dots represent the averaged values of the folded model light curve of the odd/even transits fit; the green dots are the averaged folded fit residuals, vertically offset for clarity. Odd-even transits fit completed with full convergence.

Open `./planet-01/planet-search-and-model-fitting-results/odd-even-transits-fit/0000000283722336-01-odd-even-whitened.fig`




Folded flux time series for CatId 283722336, Planet candidate 1 in the whitened domain, zoomed on the transit. The flux data whose robust weights are larger/smaller than 0.1 are plotted in dark green/cyan dots, respectively. Values are averaged into 1 cadence wide bins. The blue dots represent the averaged values of the folded flux time series; the red dots represent the averaged values of the fitted model light curve of the odd/even transits fit; the green dots are the averaged folded fit residuals, vertically offset for clarity. Magenta dots are the averaged values of the folded flux time series, with a phase shift of 0.5 relative to the blue dots, vertically offset for clarity. Odd-even transits fit completed with full convergence.

Open `./planet-01/planet-search-and-model-fitting-results/odd-even-transits-fit/0000000283722336-01-odd-even-whitened-zoomed.fig`




Robust weights distribution for CatId 283722336, Planet candidate 1. Top plot: all data points. Middle plot: all data points, folded per the fitted period and epoch. Bottom plot: all data points, folded and zoomed.

Open `./planet-01/planet-search-and-model-fitting-results/odd-even-transits-fit/0000000283722336-01-odd-even-robust-weights.fig`



Fit residuals distribution for CatId 283722336, Planet candidate 1. Only the valid data points used to constrain the fit are shown here. A Gaussian fit to the histogram is shown in red.



Open `./planet-01/planet-search-and-model-fitting-results/odd-even-transits-fit/0000000283722336-01-odd-even-histo-used.fig`



Fit residuals distribution for CatId 283722336, Planet candidate 1. Top plot: all valid data. Bottom plot: valid data not used to constrain fit (due to distance from a transit). Gaussian fits to the histograms are shown in red.

Open `./planet-01/planet-search-and-model-fitting-results/odd-even-transits-fit/0000000283722336-01-odd-even-histo-all-and-unused.fig`

### A.3 Eclipsing Binary Discrimination Test




Top-left: Diagnostic plot of Odd/Even Transit Depth Test for catId 283722336, planet 1. A significance level close to 1/0 favors a transiting planet/an eclipsing binary.  
 Bottom-left: Diagnostic plot of Orbital Period Test for catId 283722336. Orbital periods of planet 1 and the planet with longer period are compared. A significance level close to 1/0 favors a transiting planet/an eclipsing binary.

Open `./planet-01/binary-discrimination-test-results/0000000283722336-01-eclipsing-binary-discrimination-tests.fig`


## Appendix B Planet Candidate 2

### B.1 Model Fitter: All Transits




Robust weights distribution for CatId 283722336, Planet candidate 2. Top plot: all data points. Middle plot: all data points, folded per the fitted period and epoch. Bottom plot: all data points, folded and zoomed.

Open `./planet-02/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-02-all-robust-weights.fig`



Fit residuals distribution for CatId 283722336, Planet candidate 2. Only the valid data points used to constrain the fit are shown here. A Gaussian fit to the histogram is shown in red.

Open `./planet-02/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-02-all-histo-used.fig`



Fit residuals distribution for CatId 283722336, Planet candidate 2. Top plot: all valid data. Bottom plot: valid data not used to constrain fit (due to distance from a transit). Gaussian fits to the histograms are shown in red.


Open `./planet-02/planet-search-and-model-fitting-results/all-transits-fit/0000000283722336-02-all-histo-all-and-unused.fig`



## B.2 Model Fitter: Odd &amp; Even Transits


Parameter	Odd Transits Value	Odd Transits Uncertainty	Even Transits Value	Even Transits Uncertainty	Units	$\frac{\text{Difference}}{\ \text{Uncertainty}\ }$
SNR	12.1		12.2			
Orbital Period	6.7660167	9.7999e-04	6.7650130	1.0394e-03	days	7.0262e-01
Transit Epoch	1766.1687683	1.4776e-03	1772.9349988	1.5009e-03	BTJD	3.7814e-01
Impact Parameter	0.7631	9.4181e-01	0.8971	1.3362e-01		1.4079e-01
Planet Radius to Star Radius Ratio	0.0173696	6.5436e-03	0.0179923	2.3232e-03		8.9681e-02
Semi-major Axis to Star Radius Ratio	21.9298	3.7152e+01	15.1245	9.0762e+00		1.7794e-01
Planet Radius	1.4233	5.4719e-01	1.4743	2.2138e-01	Earth radii	8.6447e-02
Semi-major Axis	0.0650	5.9567e-03	0.0650	5.9561e-03	AU	7.6349e-04
Effective Stellar Flux	67.9388	1.3488e+01	67.9523	1.3490e+01	Goldilocks	7.0456e-04
Equilibrium Temperature	732	3.6342e+01	732	3.6344e+01	Kelvin	7.0456e-04
Stellar Density	3.0951	1.5731e+01	1.0156	1.8285e+00	Solar density	1.3131e-01
Transit Depth	300	2.3468e+01	274	2.1797e+01	ppm	8.0956e-01
Transit Duration	1.5870	2.0314e-01	1.6475	1.6474e-01	hours	2.3152e-01
Transit Ingress Duration	0.0635	2.4102e-01	0.1399	1.8888e-01	hours	2.4972e-01
Eccentricity	0.0000	0.0000e+00	0.0000	0.0000e+00		
Peri Longitude	0.0000	0.0000e+00	0.0000	0.0000e+00	degrees	
Model Chi Square Statistic (DoF)	684.3 (917.8)		684.3 (917.8)			

DoF: Degrees of Freedom




Folded flux time series for CatId 283722336, Planet candidate 2 in the whitened domain is plotted in black dots. Values are averaged into 1 cadence wide bins. The blue dots represent the averaged values of the folded flux time series; the red dots represent the averaged values of the folded model light curve of the odd/even transits fit; the green dots are the averaged folded fit residuals, vertically offset for clarity. Odd-even transits fit completed with full convergence.

Open `./planet-02/planet-search-and-model-fitting-results/odd-even-transits-fit/0000000283722336-02-odd-even-whitened.fig`




Folded flux time series for CatId 283722336, Planet candidate 2 in the whitened domain, zoomed on the transit. The flux data whose robust weights are larger/smaller than 0.1 are plotted in dark green/cyan dots, respectively. Values are averaged into 1 cadence wide bins. The blue dots represent the averaged values of the folded flux time series; the red dots represent the averaged values of the fitted model light curve of the odd/even transits fit; the green dots are the averaged folded fit residuals, vertically offset for clarity. Magenta dots are the averaged values of the folded flux time series, with a phase shift of 0.5 relative to the blue dots, vertically offset for clarity. Odd-even transits fit completed with full convergence.

Open `./planet-02/planet-search-and-model-fitting-results/odd-even-transits-fit/0000000283722336-02-odd-even-whitened-zoomed.fig`




Robust weights distribution for CatId 283722336, Planet candidate 2. Top plot: all data points. Middle plot: all data points, folded per the fitted period and epoch. Bottom plot: all data points, folded and zoomed.

Open `./planet-02/planet-search-and-model-fitting-results/odd-even-transits-fit/0000000283722336-02-odd-even-robust-weights.fig`



Fit residuals distribution for CatId 283722336, Planet candidate 2. Only the valid data points used to constrain the fit are shown here. A Gaussian fit to the histogram is shown in red.



Open `./planet-02/planet-search-and-model-fitting-results/odd-even-transits-fit/0000000283722336-02-odd-even-histo-used.fig`



Fit residuals distribution for CatId 283722336, Planet candidate 2. Top plot: all valid data. Bottom plot: valid data not used to constrain fit (due to distance from a transit). Gaussian fits to the histograms are shown in red.

Open `./planet-02/planet-search-and-model-fitting-results/odd-even-transits-fit/0000000283722336-02-odd-even-histo-all-and-unused.fig`

### B.3 Eclipsing Binary Discrimination Test



Top-left: Diagnostic plot of Odd/Even Transit Depth Test for catId 283722336, planet 2. A significance level close to 1/0 favors a transiting planet/an eclipsing binary.  
 Top-right: Diagnostic plot of Orbital Period Test for catId 283722336. Orbital periods of planet 2 and the planet with shorter period are compared. A significance level close to 1/0 favors a transiting planet/an eclipsing binary.

Open `./planet-02/binary-discrimination-test-results/0000000283722336-02-eclipsing-binary-discrimination-tests.fig`

## Appendix C Alerts

Time	Severity	Message
1802.6382	warning	ghost diagnostic test will not be performed for saturated target (target=1, catId=283722336, component=ghostDiagnosticTests)